

ELECTROMAGNETIC SIGNALS FROM PLANETARY COLLISIONS

BING ZHANG & STEINN SIGURDSSON

Department of Astronomy & Astrophysics, Pennsylvania State University, University Park, PA 16802

Accepted for publication in ApJ Letters

ABSTRACT

Planet–planet collisions are expected during the early stages of the formation of extra-solar planets, and are also possible in mature planetary systems through secular planet–planet perturbations. We investigate the electromagnetic signals accompanied with these planetary collisions and their event rate, and explore the possibility of directly detecting such events. A typical Earth–Jupiter collision would give rise to a prompt EUV-soft-X-ray flash lasting for hours and a bright IR afterglow lasting for thousands of years. It is suggested that with the current and forthcoming observational technology and facilities, some of these collisional flashes or the post-collision remnants could be discovered.

Subject headings: Planetary systems

1. INTRODUCTION

More than 100 extra-solar planets have been detected. The growing sample allows us to study the properties of these extra-solar planetary systems, and sheds light on the origin of our own solar system. At the same time, our understanding of astrophysical phenomena has been greatly boosted through studying cataclysmic transient events such as supernovae, X-ray bursts, gamma-ray bursts and their afterglows. Here we discuss another type of electromagnetic transient events that arises from collisions of extra-solar planets. Such giant impacts have been long predicted within the theories of planet formation (e.g. Wetherill 1990). Giant impacts have been invoked to interpret the formation of the Earth–Moon (Wetherill 1985; Benz, Slattery & Cameron 1986; Canup & Asphaug 2001) and the Pluto–Charon (Stern 1992 and references therein) systems, as well as the anomalous obliquities of Uranus and some other solar giant planets (e.g. Slattery, Benz & Cameron 1992). Collisions are also expected in mature planetary systems due to slow onset, secular dynamical instabilities, although these collisions are less frequent than in dynamically young systems. The probability of collisions is enhanced in white dwarf planetary systems because of the perturbation caused by drastic mass loss during the formation of the white dwarfs (Debes & Sigurdsson 2002). Accompanying an impact event, a prompt transient electromagnetic signal and a fading afterglow are expected, but these signals have so far been sparsely studied. Even before the discovery of planetary systems around main sequence stars, Stern (1994) discussed the frequency and infrared (IR) signals of planetary collisions. In this Letter, we will sketch the likely electromagnetic signals from planetary collisions (§2). We will estimate the possibility of directly detecting such collisional events and their remnants, and discuss the observational strategies (§3).

2. EUV-SOFT-X-RAY FLASHES AND IR AFTERGLOWS

As an example, we consider a Jovian planet (primary) with mass $M_1 = 10^{30} \text{ g } m_J$ and radius $R_1 = 7.0 \times 10^9 \text{ cm } r_J$ being hit by a terrestrial planet (secondary) with

mass $M_2 = 6 \times 10^{27} \text{ g } m_\oplus$ and radius $R_2 = 6.4 \times 10^8 \text{ cm } r_\oplus$, where m_J and r_J are the mass and radius in units of the Jupiter values, while $m_\oplus$ and $r_\oplus$ are the mass and radius in units of the Earth values, respectively. The reasons to choose a Jupiter-sized object being the primary are several folds. (1) Over 100 extrasolar giant planets (EGPs) have been detected, so that we know they are definitely there; (2) The signal is the strongest for these impacts because of their large masses (and hence higher Eddington luminosities); (3) Because of their larger gravitational potential and larger cross-sectional area, the chance of collision per planet is higher at higher masses. Since the main emission quantities scale with masses and radii, the discussion below can be straightforwardly generalized to collisions between planets with other parameters (e.g. EGPs).

The probability of a glancing event is $\sim R_2/R_1$ for a gravitational-focused system, or $\sim (R_2/R_1)^2$ when the colliding velocity largely exceeds the escaping velocity of the primary. For our parameters, the case is marginal for both regimes, and the probability for glancing is between $\sim 9\%(r_\oplus/r_J)$ and $\sim 1\%(r_\oplus/r_J)^2$, which is negligible. Hereafter we will assume head-on collisions throughout¹. Assuming zero velocity at infinity, the total energy of the collision is $E \sim GM_2M_1/R_1 \sim 6 \times 10^{40} \text{ erg } (m_\oplus m_J/r_J)$, and the relative velocity upon impact is $v \sim 45 \text{ km s}^{-1} (m_J/r_J)^{1/2}$. Earthquake data reveal that the average p-wave speed interior of the earth is $\sim 10 \text{ km s}^{-1}$ (Dziewonski & Anderson 1981), so we can generally define a sound speed $C_s \sim 10 \text{ km s}^{-1} c_s$, with $c_s \sim 1$ for terrestrial planets. Since $v \gg C_s$, strong shocks will propagate into both colliders, converting most of the kinetic energy into internal energy and accelerating non-thermal electrons. The shock crossing time for the impactor, i.e., the terrestrial planet in our case, is

$$\tau_{sh} \sim R_2/C_s \sim 640 \text{ s } (r_\oplus/c_s). \quad (1)$$

This defines the rising timescale of the collision-induced electromagnetic flash. The energy deposit rate during this stage is $\dot{E} \sim E/\tau_{sh} \sim 9 \times 10^{37} \text{ erg s}^{-1} (m_\oplus m_J/r_J)(r_\oplus/c_s)$.

¹We note that for collisions between planets with similar masses, the chance of glancing events is much larger, which leads to warm planets rather than the signals discussed in this Letter.

This is $\sim 1.8 \times 10^3$ times the Eddington luminosity of the Jovian planet, $L_{\text{Edd}} = 4\pi c G m_p M / \sigma_T \sim 5 \times 10^{34} \text{ erg s}^{-1} m_J$ (where c , G , m_p , σ_T are the speed of light, gravitational constant, proton mass, and Thomson cross section, respectively). About one half of the collisional energy is converted to internal energy, of which only a small fraction is radiated promptly, while a large fraction is sunk as latent heat. The other half of the energy is initially in kinetic form, driving a radiation-supported expanding envelope and inducing oscillation and convection within the giant planet interior. This kinetic energy will be thermalized later and radiated away over a longer period of time along with the latent heat, as an IR afterglow (see below). In the prompt phase, photons must be trapped by the high opacity in the regions where the heat is initially deposited, and the peak bolometric luminosity is

$$L_{pk} = \eta L_{\text{Edd}} \sim 5 \times 10^{34} \text{ erg s}^{-1} \eta m_J \sim 13 L_{\odot} \eta m_J, \quad (2)$$

where $L_{\odot} = 3.8 \times 10^{33} \text{ erg s}^{-1}$ is the solar luminosity, $\eta \leq 1$ is a factor correcting for less than optimal emission geometry and other radiative inefficiencies. Since $\dot{E} \gg L_{\text{Edd}}$, $\eta \sim 1$ is likely achieved. The spectrum is essentially thermal from a hot spot² with radius $\sim R_2$, with a non-thermal hard tail caused by Comptonization of the thermal photons by the non-thermal electrons accelerated from the shocks. The typical thermal temperature at the peak time is

$$T_{pk} = \left(\frac{L_{pk}}{4\pi R_2^2 \sigma} \right)^{1/4} \sim 1.1 \times 10^5 \text{ K } \eta^{1/4} m_J^{1/4} r_{\oplus}^{-1/2}, \quad (3)$$

which is insensitive to both the planet mass and the unknown radiation efficiency (i.e. $\propto \eta^{1/4} m_J^{1/4}$). The peak frequency is $\nu_{pk} = 6.6 \times 10^{15} \text{ Hz } \eta^{1/4} m_J^{1/4} r_{\oplus}^{-1/2}$ (wavelength $\sim 450 \text{ \AA}$ and energy $\sim 28 \text{ eV}$). This is in the extreme ultraviolet (EUV) band. The peak flux is

$$F_{\nu}(pk) = \frac{h\sigma T^3}{2.8k} \cdot \left(\frac{R_2}{D} \right)^2 = 60 \mu\text{Jy } \eta^{3/4} m_J^{3/4} r_{\oplus}^{1/2} \left(\frac{D}{10\text{kpc}} \right)^{-2}, \quad (4)$$

where h , k , and σ are the Planck's, Boltzmann's and Stefan-Boltzmann constants, respectively. The hard non-thermal tail is expected to extend into the soft-X-ray band.

For the thermal spectrum, the flux around ν could be estimated as $F(\nu) \sim (R/D)^2 B_{\nu}(T)\nu$, where $B_{\nu}(T)$ is the Planck function, R is the emission radius, and D is the distance to Earth. One can define a ratio $f(\nu) \equiv F(\nu, p)/F(\nu, *)$ as the flux contrast between the planet and the star at a certain frequency ν , where $F(\nu, p)$ and $F(\nu, *)$ are the fluxes of the planet and the star in the band around ν , respectively. At the flash peak time, the EUV flare can greatly out-shine a main-sequence host star later than B5, or a white dwarf host star. In the optical and IR bands, high contrasts are also achievable for late-type host stars. For example, for a G2 (solar-type) star, the U, V, and I bands contrasts at the peak time are $f(\text{U}, pk) \sim 0.2$, $f(\text{V}, pk) \sim 0.02$, and $f(\text{I}, pk) \sim 0.008$, respectively. Even higher contrasts are achievable when a cooler star is adopted. For $f \geq 0.01$, the flare is detectable

with photometric monitoring of ground-based telescopes (e.g. Borucki et al. 2001).

Fixing a particular band, a lightcurve could be depicted. The rising lightcurve is very steep, with a time scale of $\sim \tau_{sh} \sim 10$ minutes. The decay time scale after the peak is likely not very long, based on existing observational data of other related phenomena, e.g. Comet Shoemaker-Levy 9 vs. Jupiter collision (Orton et al. 1995), X-ray bursts in accreting neutron stars (Lewin, van Paradijs & Taam 1993) and soft gamma-ray giant flares from magnetars (Feroi et al. 2001). The prompt radiation pressure would drive a fraction of the impacting materials outwards, so that initially the Eddington-limited photosphere expands and then cools. As the radiation pressure gradually drops, the photosphere contracts and the materials falls back onto the Jovian planet surface. This fallback refreshes the surface heating, leading to continually refreshed electromagnetic flashes with descending magnitudes. A best guess is that the prompt flash will follow a decaying tail extending up to several 10's of the rising time scale. The typical timescale for the prompt flash during which the emission luminosity is above, say 50% of L_{pk} could be estimated

$$\tau_1 \sim 10\tau_{sh} \sim 2 \text{ hr } (r_{\oplus}/c_s), \quad (5)$$

with the averaged bolometric luminosity $\bar{L}_{bol}^{flash} \sim 10 L_{\odot} \eta m_J$. This is the first important time scale in the problem.

The total radiated energy during this prompt flash stage is $E^{flash} \sim \bar{L}_{bol}^{flash} \tau_1 \sim 3 \times 10^{38} \text{ erg}$, which is much less than the total impact energy, i.e. $\ll E \sim 6 \times 10^{40} \text{ ergs}$. So the bulk of the collision-deposited energy is *not* radiated promptly. Rather, it is mixed into deep inside of the planet and stored as heat. It is released during a much longer time scale through a decaying thermal-radiation afterglow. Model simulations reveal that at later times a Jovian planet follows an empirical bolometric luminosity law (Black 1980; Saumon et al. 1996), i.e., $L/L_{\odot} \sim 10^{-5} m_J^{2.35} t_6^{-1.22}$, and $T_{\text{eff}} \sim 860 \text{ K } m_J^{0.51} t_6^{-0.28}$, where t_6 is the age in unit of Myr, and the relations are valid for $t_6 > 1$. Notice that the luminosity decaying index $\sim 1.22 > 1$, so that the total energy released during $t > 10^6 \text{ yr}$ is negligible compared with the energy released during the early epoch. At earlier times, generally we expect a luminosity decaying law $L = L_0(t/t_0)^{-a}$ with $a < 1$, where t_0 and L_0 are the time and luminosity when such a steady fading phase begins. The duration of the bright afterglow (the second timescale in the problem) can be estimated as $\tau_2 = \tau_{ag} = [(1-a)E/L_0 t_0]^{1/(1-a)} t_0$. The values of t_0 , L_0 and a , are unknown, and a may also change during the decay. In principle, these should be calculated from the giant planet models. As a crude estimate, we take $(t_0, L_0) = (1 \text{ yr}, 3.6 \times 10^{-4} L_{\odot})$. The L_0 value corresponds to an initial temperature of $T_0 \sim 2500 \text{ K}$ (e.g. Stern 1994 and references therein) emitting from the full giant planet surface with the normalized radius $r_J = 1$. Given such an initial condition, $a < 1/4$ is required in order to have $L(t_6 = 1) > 10^{-5} L_{\odot}$. The bright afterglow duration

²The hot spot will eventually expand via a subsonic excavation flow (Melosh 1989) or a convective flow, but the timescale for this to happen is ~ 1 day, much longer than the flash timescale discussed here (eq.[5]).

is therefore

$$1.4 \times 10^3 \text{ yr } (m_{\oplus} m_J) r_J^{-3} \left(\frac{T_0}{2500 \text{ K}} \right)^{-4} \lesssim \tau_2 \lesssim 10^4 \text{ yr}, \quad (6)$$

where the lower and the higher limits correspond to $a = 0$ and $a = 1/4$, respectively, and for the lower limit, the explicit dependences on the model parameters have been introduced. The peak of this bright afterglow is in the IR band, and it is more favorably discovered in longer wavelengths. For example, for a G2 host star, the typical I- and K-band planet-to-star flux contrasts are $f(\text{I, ag}) \sim 2.6 \times 10^{-4}$ and $f(\text{K, ag}) \sim 1.7 \times 10^{-3}$ (see also Stern 1994).

Between the prompt flash and the onset of the later shallow-decay long-term afterglow, the bolometric luminosity has to drop by 5 orders of magnitude with a decay index $a > 1$. This would nicely match the (t_0, L_0) initial condition adopted in the above discussion. In the above estimate we have assumed that the prompt flash radiation is released from the hot spot, while the long-term afterglow emission is emitted from the full surface. It is inferred that the Jupiter interior must be convective (Hubbard 1968; Bishop & DeMarcus 1970). Thus the collision-deposited energy should be rapidly transported to the planet bulk during tens or hundreds of the convective overturn time scale. The depth into which the impactor velocity is reduced to $1/2$ is $\sim 2.5 \times 10^9 \text{ cm}$, i.e., the depth where the impacted mass equal to the mass of the impactor, which is about $1/3$ of the Jovian planet radius³. This length scale may be adopted as the convective mixing-length. The convective speed is less clear due to the unknown interior viscosity. Nonetheless it should be subsonic, and we take $v_c \sim 1 \text{ km s}^{-1} v_{c,0}$ for the following discussions. This gives a convective overturn timescale $\tau_{conv} \sim 0.3 v_{c,0}^{-1}$ days. The timescale for the excavative flows (Melosh 1989), if any, is longer than this. Taking a typical overturn number of $N_c = 100 N_{c,2}$ to achieve a full-surface isothermal state, we estimate a third timescale

$$\tau_3 \sim 1 \text{ month } N_{c,2} v_{c,0}^{-1}. \quad (7)$$

Since the typical spin period of giant planets is $P_p < 1 \text{ day} \ll \tau_3$, one expects a periodic luminosity modulation on the EUV-Soft-X-ray lightcurve, caused by the expanded hot spot (about 3% of the full surface) entering and leaving the line-of-sight, over a time scale of τ_3 . This may be also detectable in the U, B bands or even in the V band through photometry.

During the vigorous convective epoch, it is likely that the convection-induced turbulence would amplify the magnetic field to higher values than conventional (e.g. sub-Gauss for Jupiter). The fast relativistic electrons accelerated from the shocks or from magnetic reconnections would radiate coherently, giving rise to a radio flare similar to the Type-II solar radio flares (but with much higher amplitudes). For a flare with brightness temperature $T_b > 10^{15} \text{ K}$, the flux would out-shine the host star, and is

$\sim 1 \mu\text{Jy}$ at a distance of 10 kpc. Only very nearby events (e.g. within 1 kpc) are favorable for radio observations.

3. DETECTABILITY AND SEARCHING STRATEGY

Numerical simulations (Chambers, Wetherill & Boss 1996; Ford, Havlickova & Rasio 2001) suggest that collisions of the type we are discussing are plausible. The basic picture is that there are secular perturbations of the inner planets, which over time scales comparable to the age of the system (Quinlan & Tremaine 1992) lead to large changes in eccentricity and semi-major axis for one or more planets, leading to a large probability of collision. This is even favorable for those EGPs that are found much closer to the host star than Jupiter. Current theories conjecture that these EGPs migrate towards the host star within the gas disk from which they were born (Lin, Bodenheimer & Richardson 1996). The disk disappears in about 10 million years. Recent meteorite isotope studies suggest that terrestrial planets form in a mean time scale of 10 million years (Yin et al. 2002; Kleine et al. 2002; Jacobsen 2003), so that they could be available for the collisions we conjecture⁴.

Considering an ensemble of stars with an average age $\bar{t}$, in order to detect one event with duration τ (e.g. τ_1 for the prompt flash and τ_2 for the afterglow) after a continuous observation time of t_{obs} , the critical number of stars in this ensemble that have to be searched is

$$N_* = (f_p \bar{N}_c)^{-1} \frac{\bar{t}}{\max(\tau, t_{obs})}, \quad (8)$$

where f_p is the fraction of the stars in the ensemble that have planets, and $\bar{N}_c$ is the average total number of collisions during the lifetime of a typical star in the ensemble. Evidence suggests that f_p could be close to unity (e.g. Lineweaver & Grether 2003; Bary, Weintraub & Kastner 2002), and we will assume this most optimistic case. $\bar{N}_c$ could be roughly estimated as the number of planets in the dynamical system (which is greater at earlier epochs since each dynamical interaction destroys one planet) multiplied by the probability of direct collisions (rather than dynamical ejections), which is in the 40% to 80% range (Ford et al. 2001). The high eccentricity of some EGP orbits may be caused by dynamical ejection of another EGP. Although no collision is involved in such events, the eccentric orbits they result more likely lead to strong secular perturbations of the remaining planets on a longer time scale, particularly so if we assume inclinations are increased as well. These actually increase the probability of future collisions with planets that might otherwise be unperturbed. A good portion of collisions should happen in dynamically young systems (which associate with star forming regions), but there could be also some collisions occurring in mature planetary systems. For an old planetary system like our Solar system, a typical value of $\bar{N}_c \sim 5$ may be reasonable⁵, while for dynamically young systems,

³The penetration depth at which most of energy is deposited is even deeper, especially when the mantle of the terrestrial planet is stripped (Canup & Asphaug 2001), but not much larger than the value quoted here.

⁴In case of a late formation epoch for terrestrial planets (Wetherill 1985), a proto-terrestrial planet gives a smaller total impact energy and a shorter IR afterglow, but the prompt, Eddington-limited, flash luminosity essentially remains the same (eq.[2]).

⁵Although the lack of a major planet in the asteroid belt and the relatively small mass of Mars may be attributed to Jupiter's perturbations that inhibited the growth of a terrestrial planet near Jupiter required for collision, we note that the solar system might be atypically stable (as compared with other extrasolar systems discovered) for anthropic reasons.

one may adopt $\bar{N}_c \sim 3$. A large fraction of stars are in binaries or multiples. Many of these are either detached or very compact, allowing for the presence of planets orbiting one or both stars (Holman & Wiegert 1999; Hatzes et al 2003). Planets may be less common around stellar binaries, but those systems may contribute disproportionately to the net collision rate, as a stellar companion can drive long term secular instabilities among multiple planet systems, increasing the collision rate per system.

Besides N_* , one can also define a characteristic flux $f_{\nu,c}$ of the collisional events. Given a number density n_* of the stars with the average age $\bar{t}$, one can estimate a critical distance one has to search in order to find one collisional event, i.e., $D_c \sim (N_*/n_*)^{1/3}$. The typical flux can be then estimated with D_c . Although N_* is sensitive to the average age $\bar{t}$ of the ensemble of stars being investigated, D_c and $f_{\nu,c}$ are essentially independent of the ensemble adopted, given a constant birth rate of stars.

For the prompt flashes, since there are considerably fewer point sources in the EUV and soft X-ray sky and since the collisional events out-shine the host stars, one does not need to monitor the individual stars in order to identify a collisional event. The characteristic flux $f_{\nu,c}$ is relevant to discuss the detectability. The *Extreme Ultraviolet Explorer (EUVE)* performed a one-year EUV all-sky survey (Bowyer et al. 1994) and subsequently conducted the serendipitous observations with the Right Angle Program (RAP) (McDonald et al. 1994; Christian et al. 1999; Christian 2002) for 8 more years. The RAP field of view is ~ 20 square degrees. So the combined observation is equivalent to a one-year monitoring with $9 \times (20/720) = 1/4$ full-sky coverage. The total number of stars with $\bar{t} \sim 10^{10}$ yr, for which one collisional event is expected to have occurred in the *EUVE* field of view, is $\sim 8 \times 10^9$. The typical distance of such an event is $\sim 10 - 20$ kpc, so that the critical peak flux is $f_{\nu,c} \sim (15 - 60)\mu\text{Jy}$ peaking around $450\text{\AA}$. This is the band for which the ISM opacity is the largest. At shorter wavelengths where opacity is much lower, e.g., $50-100\text{\AA}$ ($\sim 0.1 - 0.2$ keV), the flux does not drop rapidly due to the non-thermal tail. Even for a flux two orders of magnitude lower than the peak flux (a very conservative estimate), i.e., $(0.15 - 0.6)\mu\text{Jy}$, this corresponds to $(170 - 680)$ counts ks^{-1} at $100\text{\AA}$ for *EUVE* (Bowyer et al. 1994), which is well above the survey sensitivity threshold (McDonald et al. 1994). So at least one flare event with such a flux level should have been detected. Since during ~ 10 yrs, $\sim 10 \times 10^{11}/(10^{10}/5) \sim 500$ collisional events should have occurred in the Milkyway Galaxy (~ 50 per year), about 10 more events with lower flux levels might have also been caught by *EUVE*, although with less significance. The predicted flux level is also well above the sensitivity of soft X-ray detectors, such as *ROSAT*, *Chandra*, and *XMM-Newton*, so that there might be such events recorded in their archival data as well. A future dedicated wide field detector sensitive to $(50-200)\text{\AA}$ would be able to detect 10's of EUV-soft-X-ray flares per year. The great opacity in the star forming regions would reduce the flux level and even the detectable event rate in the softer bands, but some events should be directly detected, especially for those collisions outside of the star forming regions.

Another approach to identify prompt flashes is through photometrically monitoring a huge number of stars over a

long period of time. As discussed above, during the flare, a 2-20% percent increase of the fluxes are expected for the late type stars in the V band and bluer, which could be easily detected. Such a photometric search could be achieved by future missions such as *GAIA* (e.g. Perryman et al. 2001). Close monitoring $\sim 10^9$ stars for several years should lead to detection of several collisional events.

The event rate of the IR afterglows is much larger thanks to their much longer durations. Due to small flux contrasts ($\sim 10^{-3}$ in K-band), one has to monitor the stars individually. The most economic way is to search in the star forming regions (Stern 1994). For an ensemble of stars with $\bar{t} \sim 5 \times 10^7$ yr, one IR afterglow remnant would be found by searching for $N_*^{ag} \sim 1.7 \times 10^3 f_p^{-1} (\bar{N}_e/3)^{-1} (\bar{t}/5 \times 10^7) \tau_{2,4}^{-1}$ young stars, where $\tau_{2,4} = \tau_2/10^4$ yr. The signals could be identified through unresolved spectral measurements or direct interferometric imaging (Stern 1994). Existing groundbased IR interferometers have 1 mas resolutions (J. Ge, personal communication), which corresponds to 0.13 AU at a distance of 130 pc, the distance of the nearest star forming region. This is well adequate to resolve bright planets from their host stars.

4. SUMMARY AND DISCUSSION

We have outlined the electromagnetic signals that would accompany with an event of collision between a Jovian planet and a terrestrial planet, and discussed the strategy of possibly detecting these events. The collision itself results in a bright EUV-soft-X-ray flare lasting for hours. *EUVE* and some soft X-ray detectors might have detected some of these flares. They may be also identified through monitoring a huge number of stars with future missions such as *GAIA*, or through future synoptic all-sky surveys. After detecting an EUV-soft-X-ray flare, a search of its optical counterpart through photometric monitoring (like catching afterglows in gamma-ray burst study) is desirable. A planetary-rotation-period modulated fading signal would be an important clue. Doppler radial velocity measurements and IR monitoring may be performed later for the collision candidate to verify the existence of the planet(s). Each collisional event would leave a bright remnant glowing in IR. The duration of the afterglow is thousands of years. Such afterglows could be directly searched in the nearby star forming regions, and the planets could be directly imaged. The current planet-searching has been focused on the planets on stable orbits (Perryman 2000). Searching for planetary collisional events will open a new window in the planetary science. It allows us to witness the gigantic events that are believed to have happened in the early history of our solar system, and to directly test the planet-moon-formation theories (see also Stern 1994).

We are grateful to Alan Stern, Damian Christian, Jian Ge, David Weintraub and Kai Cai for helpful comments and discussions, and to the referee for insightful comments that lead to improvement of the manuscript. B.Z. acknowledges NASA NAG5-9192 for support. S.S. is supported by NSF grant PHY-0203046, the Center for Gravitational Wave Physics (which is supported by the NSF under cooperative agreement PHY 01-14375), and the Penn State Astrobiology Research Center.

REFERENCES

- Bary, J. S. & Weintraub, D. A. & Kastner, J. H. 2002, *ApJ*, 576, L73
 Bishop, E. V. & DeMarcus, W. C. 1970, *Icarus*, 12, 317
 Benz, W., Slattery, W. L. & Cameron, A. G. W. 1986, *Icarus*, 66, 518
 Black, D. C. 1980, *Icarus*, 43, 293
 Borucki, W. J. et al. 2001, *PASP*, 113, 439
 Bowyer, S. et al. 1994, *ApJS*, 93, 569
 Canup, R. M. & Asphaug, E. 2001, *Nature*, 412, 708
 Chambers, J. E., Wetherill, G. W., & Boss, A. P. 1996, *Icarus*, 119, 261
 Christian, D. J. 2002, *AJ*, 124, 3478
 Christian, D. J. et al. 1999, *AJ*, 117, 2466
 Debes, J. H. & Sigurdsson, S. 2002, *ApJ*, 572, 556
 Dziewonski A. M. & Anderson D. L. 1981, *Phys. Earth Planet. Inter.* 25, 297
 Feroci, M., Hurley, K., Duncan, R. C. & Thompson, C. 2001, *ApJ*, 549, 1021
 Ford, E. B., Havlickova, M. & Rasio, F. A. 2001, *Icarus*, 150, 303
 Hatzes, A.P., 2003, *ApJ*, submitted (astro-ph/0305110)
 Holman, M.J. & Wiegert, P.A., 1999, *AJ*, 117, 621
 Hubbard, W. B. 1968, 152, 745
 Jacobsen, S. B. 2003, *Science*, 300, 1513
 Kleine, T. et al. 2002, *Nature*, 418, 952
 Lewin, W. H. G., van Paradijs, J. & Taam, R. E. 1993, *Space Sci. Rev.* 62, 223
 Lin, D. N. C., Bodenheimer, P. & Richardson, D. C. 1996, *Nature*, 380, 606
 Lineweaver, C.H. & Grether, D., 2003, preprint (astro-ph/0306524)
 Melosh, H. J. 1989, *Impact Cratering: A Geologic Process*, Oxford University Press
 McDonald, K. et al. 1994, *AJ*, 108, 1843
 Orton, G. et al. 1995, *Science*, 267, 1277
 Perryman, M. A. C. 2000, *Rep. Prog. Phys.* 63, 1209
 Perryman, M. A. C., et al. 2001, *A&A*, 369, 339
 Quinlan, G. D. & Tremaine, S. 1992, *MNRAS*, 259, 505
 Saumon, D., Hubbard, W. B., Burrows, A., Guillot, T., Lunine, J. I. & Chabrier, G. 1996, *ApJ*, 460, 993
 Slattery, W. L., Benz, W. & Cameron, A. G. W. 1992, *Icarus*, 99, 167
 Stern, S. A. 1992, *ARA&A*, 30, 185
 —. 1994, *AJ*, 108, 2312
 Wetherill, G. W. 1985, *Science*, 228, 877
 —. 1990, *Ann. Rev. Earth Planet. Sci.* 18, 205
 Yin, Q. et al. 2002, *Nature*, 418, 949